

Country	Recommended Chemoprophylaxis⁵
Australia	Not Applicable
Bangladesh	Atovaquone-proguanil, doxycycline, or mefloquine.
Bhutan	Atovaquone-proguanil, doxycycline, or mefloquine
Brunei	Not Applicable
Cambodia	In the provinces of Banteay Meanchey, Battambang, Kampot, Koh Kong, Odder Meanchey, Pailin, Preah Vihear, Pursat, and Siem Reap bordering Thailand: Atovaquone-proguanil or doxycycline All other areas with malaria: Atovaquone-proguanil, doxycycline, or mefloquine Phnom Penh: mosquito avoidance only
China	Along China-Burma (Myanmar) border in the western part of Yunnan province: Atovaquone-proguanil or doxycycline Hainan and other parts of Yunnan province: Atovaquone-proguanil, doxycycline, or mefloquine Anhui, Guizhou, Henan, and Hubei provinces: Atovaquone-proguanil, chloroquine, doxycycline, mefloquine, or primaquine ⁷ All other areas with malaria including river cruises that pass through malaria-endemic provinces: Mosquito avoidance only.
Cook Islands	Not Applicable
Fiji	Not Applicable
French Polynesia	Not Applicable
India	Atovaquone-proguanil, doxycycline, or mefloquine
Indonesia	Atovaquone-proguanil, doxycycline, or mefloquine
Japan	Not Applicable
Kiribati (formerly Gilbert Islands).	Not Applicable
Lao People's Democratic Republic	Along the Laos-Burma (Myanmar) border in the provinces of Bokeo and Louang Namtha and along the Laos-Thailand border in the province of Champasack and Saravan: Atovaquone-proguanil or doxycycline. All other areas with malaria: Atovaquone-proguanil, doxycycline, or mefloquine

Malaysia	Atovaquone-proguanil, doxycycline, or mefloquine
Maldives	Not Applicable
Marshall Islands	Not Applicable
Micronesia, Federated States of;	Not Applicable
Mongolia	Not Applicable
Myanmar (Burma)	In the provinces of Bago, Kachin, Kayah, Kayin, Shan, and Tanintharyi: Atovaquone-proguanil or doxycycline. All other areas with malaria: Atovaquone-proguanil, doxycycline, or mefloquine.
Nauru	Not Applicable
Nepal	Atovaquone-proguanil, doxycycline, or mefloquine
New Caledonia (France)	Not Applicable
New Zealand	Not Applicable
Niue (New Zealand)	Not Applicable
Korea, North	Atovaquone-proguanil, chloroquine, doxycycline, mefloquine, or primaquine ⁷
Pakistan	Atovaquone-proguanil, doxycycline, or mefloquine
Palau	Not Applicable
Papua New Guinea	Atovaquone-proguanil, doxycycline, or mefloquine
Philippines	Atovaquone-proguanil, doxycycline, or mefloquine
Samoa	Not Applicable
Singapore	Not Applicable
Solomon Islands	Atovaquone-proguanil, doxycycline, or mefloquine
Sri Lanka	Not Applicable
Korea, South	Atovaquone/ proguanil, chloroquine, doxycycline, mefloquine, or primaquine ⁷

Thailand	Atovaquone-proguanil or doxycycline
Timor-Leste (East Timor)	Atovaquone-proguanil, doxycycline, or mefloquine
Tokelau (New Zealand)	Not Applicable
Tonga	Not Applicable
Tuvalu	Not Applicable
Vanuatu	Atovaquone-proguanil, doxycycline, or mefloquine
Vietnam	Southern part of the country in the provinces of Dac Lac, Gia Lai, Khanh Hoa, Kon Tum, Lam Dong, Ninh Thuan, Song Be, Tay Ninh: Atovaquone-proguanil or doxycycline Other areas with malaria except Mekong Delta: Atovaquone-proguanil, doxycycline or mefloquine Mekong Delta: Mosquito avoidance

Source:

CDC (2013) Malaria Information and Prophylaxis by country (http://www.cdc.gov/malaria/travelers/country_table/a.html). The information presented in this table is consistent with the information in the **Health Information for International Travel 2014 (CDC's Yellow Book)** (<http://wwwnc.cdc.gov/travel/page/yellowbook-home-2014>).

Note:

Several medications are available for chemoprophylaxis. When deciding which drug to use, consider specific itinerary, length of trip, cost of drug, previous adverse reactions to antimalarials, drug allergies, and current medical history. All travelers should seek medical attention in the event of fever during or after return from travel to areas with malaria. All travellers should seek medical advice on choice of chemoprophylaxis.